

## Selected Dynabeads® References for Organelle Proteomics

### Electron Microscopy:

- Kumar A *et al.* Isolation and characterization of complement receptor type 1 (CR1) storage vesicles from human neutrophils using antibodies to the cytoplasmic tail of CR1. *Blood* 1997;89:4555-65.
- Hansen GH *et al.* Lipid rafts exist as stable cholesterol-independent microdomains in the brush border membrane of enterocytes. *J. Biol. Chem.* 2001;276(34):32338-32344. ([full text article - JBC Online](#))

### Cell-free assays:

- Gruenberg JE and Howell KE. Reconstitution of vesicle fusions occurring in endocytosis with a cell-free system. *The EMBO Journal* 1986;5(12):3091-3101.
- Salamero J *et al.* Exocytic transport vesicles generated in vitro from the trans-Golgi network carry secretory and plasma membrane proteins. *PNAS USA* 1990;87:7717-7721.

### Endosomes:

- Gagescu R *et al.* The recycling endosome of Madin-Darby canine kidney cells is a mildly acidic compartment rich in raft components. *Molecular Biology of the Cell* 2000;11:2775-2791.
- Li X *et al.* Na<sup>+</sup> - H<sup>+</sup> exchanger 3 (NHE3) is present in lipid rafts in the rabbit ileal brush border: a role for rafts in trafficking and rapid stimulation of NHE3. *J. Physiol.* 2001;537(2):537-552

### Golgi:

- Vitale N *et al.* Localization of ADP-ribosylation factor domain protein 1 (ARD1) in lysosomes and Golgi apparatus. *PNAS* 1998;95:8613-8618.

### Kidney glomeruli:

- Takemoto M. *et al.* (2002). A new method for large scale isolation of kidney glomeruli from mice. *Am J Pathol.* 161:799-805.

### Lens membrane fractions:

- Sexton PS and Cenedella RJ. Immunomagnetic capture of lens membrane fractions containing steroid binding protein. *Biochemical and Biophysical Research Communications* 2002;295:1027-1031. ([abstract - PubMed](#))

### Mitochondria:

- Herrnstadt C *et al.* A novel mitochondrial DNA-like sequence in the human nuclear genome. *Genomic* 1999;60:67-77

### Peroxisomes:

- Lüers GH *et al.* Immunoisolation of highly purified peroxisomes using magnetic beads and continuous immunomagnetic sorting. *Electrophoresis* 1998;19:1205-1210.

### Post-synaptic densities/synaptic vesicles:

- Sans N *et al.* Synapse-associated protein 97 selectively associates with a subset of AMPA receptors early in their biosynthetic pathway. *J. Neuroscience* 2001;21(19):7506-7516. ([full text article - Journal of Neuroscience](#))
- Provoda CJ *et al.* Evidence for a primary endocytic vesicle involved in synaptic vesicle biogenesis. *J.Biol.Chem.* 2000;275(10):7004-7012.

### Vesicles/membranes:

- Hansen GH *et al.* Lipid rafts exist as stable cholesterol-independent microdomains in the brush border membrane of enterocytes. *J. Biol. Chem.* 2001;276(34):32338-32344. ([full text article - JBC Online](#))
- Allan BB *et al.* Rab1 recruitment of p115 to a cis-SNARE complex: programming budding COPII vesicles for fusion. *Science* 2000;289(5478):444-448.
- Biemesderfer D *et al.* Membrane topology of NHE3. *J.Biol.Chem.* 1998;273(20):12391-12396.
- Calhoun BC and Goldenring JR. Two Rab proteins, vesicle-associated membrane protein 2 (VAMP-2) and secretory carrier membrane proteins (SCAMPs), are present on immunisolated parietal cell tubovesicles. *Biochem J.* 1997;325:559-564.
- Fisher MD and Frost SC. Using magnetic beads to isolate inside-Out Glut1-containing vesicles from 3T3-L1 adipocyte plasma membranes. *Anal.Biochem.* 1997;251:125-126.
- Jesch S and Linstedt AD. The Golgi and endoplasmic reticulum remain independent during mitosis in HeLa cells. *Mol.Biol. of the Cell* 1998;9:623-635.
- Lanoix J *et al.* Sorting of Golgi resident proteins into different subpopulations of COPI vesicles: a role for ArfGAP1. *J. Cell Biol.* 2001;155(7):1199-1212.
- Madey E *et al.* Characterization of plasma membrane domains enriched in lipid metabolites. *J. Exp.Bot.* 2001;52(357):669-679.
- Oh P *et al.* Dynamin at the neck of caveolae mediates their budding to form transport vesicles by GTP-driven fission from the plasma membrane endothelium. *J.Cell Biol.* 1998;141(1):101-114.
- Phillips M *et al.* The products of the *Drosophila stoned* locus interact with synaptic vesicles via synaptotagmin. *J. Neuroscience* 2000;20(22):8254-8261.
- Randhawa VK *et al.* VAMP2, but not VAMP3/cellubrevin, mediates insulin-dependent incorporation of GLUT4 into the plasma membrane of L6 myoblasts. *Mol.Biol of the Cell* 2000;11:2403-2417.
- Rehling P *et al.* Formation of AP-3 transport intermediates requires Vps41 function. *Nature Cell Biology* 1999;1:346-353.
- Safran M *et al.* Catalytic activity of type II iodothyronine 5'-deiodinase polypeptide is dependent upon a cyclic AMP activation factor. *J.Biol.Chem.* 1996;271(27):16363-16368
- Sinai AP *et al.* Association of host cell endoplasmic reticulum and mitochondria with the *Toxoplasma gondii* parasitophorous vacuole membrane: a high affinity interaction. *J.Cell Sci.* 1997;110:2117-2128
- Volchuk A *et al.* Cellubrevin is a resident protein of insulin-sensitive GLUT4 glucose transporter vesicles in 3T3-L1 adipocytes. *J.Biol.Chem.* 1995;270(14):8233-8240.